


FIG. 1A










FIG. 2

09918580.07401








FIG. 4A




FIG. 4B




FIG. 5A

FIG. 5A





FIG. 5B






FIG. 6A




FIG. 6B




FIG. 7







FIG. 8








FIG. 11B






FIG. 12A



FIG. 12B






FIG. 13A



Table 1. Demographic characteristics of the study population	
Age (years)	65.0 ± 10.0
Gender	
Male	50 (50.0%)
Female	50 (50.0%)
Education (years)	12.0 ± 2.0
Marital status	
Married	40 (80.0%)
Single	10 (20.0%)
Occupation	
Retired	30 (60.0%)
Unemployed	20 (40.0%)
Income (USD/month)	1,200 ± 300
Health status	
Good	30 (60.0%)
Fair	20 (40.0%)
Poor	10 (20.0%)
Comorbidities	
Hypertension	20 (40.0%)
Diabetes	10 (20.0%)
Cholesterol	15 (30.0%)
Arthritis	10 (20.0%)
Depression	5 (10.0%)
Other	5 (10.0%)
Medication	
Yes	30 (60.0%)
No	20 (40.0%)
Medication type	
Antidepressant	10 (20.0%)
Antipsychotic	5 (10.0%)
Anticholinergic	5 (10.0%)
Other	10 (20.0%)
Medication duration (years)	5.0 ± 3.0
Medication adherence	
High	20 (40.0%)
Low	10 (20.0%)
Medication side effects	
Yes	10 (20.0%)
No	20 (40.0%)
Side effect type	
Drowsiness	5 (10.0%)
Dry mouth	5 (10.0%)
Constipation	5 (10.0%)
Other	5 (10.0%)
Side effect duration (years)	2.0 ± 1.0
Side effect severity	
Mild	5 (10.0%)
Severe	5 (10.0%)
Side effect management	
Yes	10 (20.0%)
No	20 (40.0%)
Management type	
Medication adjustment	5 (10.0%)
Non-pharmaceutical	5 (10.0%)
Other	5 (10.0%)
Management duration (years)	1.0 ± 0.5
Management effectiveness	
High	5 (10.0%)
Low	5 (10.0%)
Management satisfaction	
Yes	10 (20.0%)
No	20 (40.0%)
Satisfaction type	
High	5 (10.0%)
Low	5 (10.0%)
Satisfaction duration (years)	1.0 ± 0.5
Satisfaction management	
Yes	10 (20.0%)
No	20 (40.0%)
Management type	
Medication adjustment	5 (10.0%)
Non-pharmaceutical	5 (10.0%)
Other	5 (10.0%)
Management duration (years)	1.0 ± 0.5
Management effectiveness	
High	5 (10.0%)
Low	5 (10.0%)
Management satisfaction	
Yes	10 (20.0%)
No	20 (40.0%)
Satisfaction type	
High	5 (10.0%)
Low	5 (10.0%)
Satisfaction duration (years)	1.0 ± 0.5
Satisfaction management	
Yes	10 (20.0%)
No	20 (40.0%)
Management type	
Medication adjustment	5 (10.0%)
Non-pharmaceutical	5 (10.0%)
Other	5 (10.0%)
Management duration (years)	1.0 ± 0.5
Management effectiveness	
High	5 (10.0%)
Low	5 (10.0%)
Management satisfaction	
Yes	10 (20.0%)
No	20 (40.0%)
Satisfaction type	
High	5 (10.0%)
Low	5 (10.0%)
Satisfaction duration (years)	1.0 ± 0.5
Satisfaction management	
Yes	10 (20.0%)
No	20 (40.0%)
Management type	
Medication adjustment	5 (10.0%)
Non-pharmaceutical	5 (10.0%)
Other	5 (10.0%)
Management duration (years)	1.0 ± 0.5
Management effectiveness	
High	5 (10.0%)
Low	5 (10.0%)
Management satisfaction	
Yes	10 (20.0%)
No	20 (40.0%)
Satisfaction type	
High	5 (10.0%)
Low	5 (10.0%)
Satisfaction duration (years)	1.0 ± 0.5
Satisfaction management	
Yes	10 (20.0%)
No	20 (40.0%)
Management type	
Medication adjustment	5 (10.0%)
Non-pharmaceutical	5 (10.0%)
Other	5 (10.0%)
Management duration (years)	1.0 ± 0.5
Management effectiveness	
High	5 (10.0%)
Low	5 (10.0%)
Management satisfaction	
Yes	10 (20.0%)
No	20 (40.0%)
Satisfaction type	
High	5 (10.0%)
Low	5 (10.0%)
Satisfaction duration (years)	1.0 ± 0.5
Satisfaction management	
Yes	10 (20.0%)
No	20 (40.0%)
Management type	
Medication adjustment	5 (10.0%)
Non-pharmaceutical	5 (10.0%)
Other	5 (10.0%)
Management duration (years)	1.0 ± 0.5
Management effectiveness	
High	5 (10.0%)
Low	5 (10.0%)
Management satisfaction	
Yes	10 (20.0%)
No	20 (40.0%)
Satisfaction type	
High	5 (10.0%)
Low	5 (10.0%)
Satisfaction duration (years)	1.0 ± 0.




FIG. 14